

REGIONAL STREAM SEDIMENT AND WATER GEOCHEMICAL RECONNAISSANCE DATA, BRITISH COLUMBIA, 1983. GSC-OF 1001, NGR 66-1983, NTS 93N

```
*****  
*                                     *  
*      OPEN FILE    1001          *  
*                                     *  
*****
```

	PAGE
SURVEY NOTES	1
DATA LIST	7
SUMMARY STATISTICS	56

OPEN FILE 1001 IS ONE OF TWO OPEN FILES 1000 AND 1001 COVERING THE CENTRAL BRITISH COLUMBIA SURVEY, COMPRISING OF NTS 93M AND 93N RESPECTIVELY.

THE RECONNAISSANCE SURVEY WAS UNDERTAKEN BY THE GEOLOGICAL SURVEY OF CANADA IN CONJUNCTION WITH THE BRITISH COLUMBIA DEPARTMENT OF ENERGY, MINES AND PETROLEUM RESOURCES UNDER THE JOINT CANADA-BRITISH COLUMBIA PROJECT.

E.H.W. HORN BROOK DIRECTED GEOLOGICAL SURVEY OF CANADA ACTIVITIES.  
W.J. MCMILLAN (B.C. GOV'T) DIRECTED B.C. DEPT. OF MINES, ENERGY AND PETROLEUM RESOURCES ACTIVITIES.

N.G. LUND AND A.C. GALLETTA WERE RESPONSIBLE FOR DATA MANAGEMENT.  
N.G. LUND CO-ORDINATED OPEN FILE PRODUCTION.  
J.YELLE SUPERVISED MAP PREPERATION.

COMPUTER AND PLOTTING FACILITIES WERE PROVIDED BY THE COMPUTER SCIENCE CENTER OF E.M.R.

CONTRACTS LET FOR SAMPLE COLLECTION, PREPARATION AND ANALYSIS WERE SUPERVISED AND/OR MONITORED BY THE STAFF OF THE GEOCHEMISTRY SUBDIVISION AND THE B.C. DEPT. AS FOLLOWS :

COLLECTION	- HARDY ASSOCIATES, BURNABY, B.C.
	- E.H.W. HORN BROOK, N.G. LUND
PREPARATION	- GOLDER ASSOCIATES, OTTAWA.
	- J.J. LYNCH
ANALYTICAL	- CHEMEX LABS. LIMITED, VANCOUVER.
	- ACME ANALYTICAL LABORATORIES LTD, TORONTO.
	- J.J. LYNCH

OPEN FILE TEXT WAS MANUFACTURED BY CAMPBELL LAZER PRINTING, OTTAWA.

HELICOPTER SUPPORTED SAMPLE COLLECTION WAS CARRIED OUT DURING THE SUMMER OF 1983.  
STREAM SEDIMENT AND WATER SAMPLES WERE COLLECTED AT AN AVERAGE DENSITY OF ONE SAMPLE PER 13 SQUARE KILOMETERS THROUGHOUT THE 28,000 SQUARE KILOMETER CENTRAL B.C.

SAMPLE SITE DUPLICATE SAMPLES WERE ROUTINELY COLLECTED IN EACH ANALYTICAL BLOCK OF TWENTY SAMPLES.

IN OTTAWA, FIELD DRIED SAMPLES WERE AIR-DRIED, CRUSHED, BALL MILLED AND SEIVED. THE MINUS 80 MESH (177 MICRONS) FRACTION WAS USED FOR SUBSEQUENT ANALYSES. AS REQUIRED, AT THIS TIME, CONTROL REFERENCE AND BLIND DUPLICATE SAMPLES WERE INSERTED INTO EACH BLOCK OF TWENTY SEDIMENT SAMPLES. NO OTHER SAMPLE PROCESSING IN OTTAWA WAS CARRIED OUT ON THE WATER SAMPLES. BLIND DUPLICATE SAMPLES WERE NOT USED IN WATER ANALYSIS.

ON RECEIPT, FIELD AND ANALYTICAL DATA WERE PUNCHED ONTO 80 COLUMN CARDS AND ALL SUBSEQUENT PROCESSING WAS CARRIED OUT WITH THE AID OF COMPUTERS. THE FIELD DATA WERE RECORDED BY THE FIELD CONTRACT STAFF ONTO STANDARD LAKE SEDIMENT FIELD CARDS (REV. 74) USED BY THE GEOLOGICAL SURVEY OF CANADA (GARRETT, 1974). THE SAMPLE SITE POSITIONS WERE MARKED ON APPROPRIATE 1/250,000 SCALE NTS MAPS IN THE FIELD. THESE MAPS WERE DIGITIZED AT THE GEOLOGICAL SURVEY IN OTTAWA TO OBTAIN THE SAMPLE SITE UTM COORDINATES.

THE SAMPLE SITE COORDINATES WERE CHECKED BY PLOTTING SAMPLING LOCATION MAPS ON A CALCOMP 1051 DRUM PLOTTER FROM THE DIGITIZED COORDINATES AND THEN OVERLAYING THESE OVER THE FIELD CONTRACTOR'S SAMPLE LOCATION BASE MAPS. THE DOMINANT ROCK TYPES IN THE LAKE CATCHMENT BASINS WERE IDENTIFIED ON APPROPRIATE GEOLOGICAL MAPS USED AS THE BEDROCK GEOLOGICAL BASE ON NGR MAPS.

THOROUGH INSPECTIONS OF THE FIELD AND ANALYTICAL DATA WERE MADE TO CHECK FOR ANY MISSING INFORMATION AND/OR GROSS ERRORS.

QUALITY CONTROL AND MONITORING OF THE GEOLOGICAL DATA WAS UNDERTAKEN BY A STANDARD METHOD USED BY THE RESOURCE GEOCHEMISTRY SUBDIVISION AT THE GEOLOGICAL SURVEY OF CANADA.

FOR THE DETERMINATION OF ZN, CU, PB, NI, CO, AG, MN AND FE, A 1 GRAM SAMPLE WAS REACTED WITH 6 ML OF A MIXTURE OF 4M HCL AND M HNO<sub>3</sub> IN A TEST-TUBE OVERNIGHT AT ROOM TEMPERATURE. AFTER DIGESTION, THE TEST-TUBE WAS IMMERSSED IN A HOT WATER BATH AT ROOM TEMPERATURE AND BROUGHT UP TO 90C AND HELD AT THIS TEMPERATURE FOR 2 HOURS WITH PERIODIC SHAKING. THE SAMPLE SOLUTION WAS THEN DILUTED TO 20 ML WITH METAL FREE WATER AND MIXED. ZN, CU, PB, NI, CO, AG, MN FE AND CD WERE DETERMINED BY ATOMIC ABSORTION SPECTROSCOPY USING AN AIR-ACETYLENE FLAME. BACKGROUND CORRECTIONS WERE MADE FOR PB, NI, CO, AG AND CD.

ARSENIC WAS DETERMINED BY ATOMIC ABSORPTION USING A HYDRIDE EVOLUTION METHOD WHEREIN THE ARSENIC IS EVOLVED AS ASH<sub>3</sub>, PASSED THROUGH A HEATED QUARTZ TUBE IN THE LIGHT PATH OF AN ATOMIC ABSORPTION SPECTROPHOTOMETER. THE METHOD IS DESCRIBED BY ASLIN (1976).

MOLYBDENUM WAS DETERMINED BY ATOMIC ABSORPTION SPECTROSCOPY USING A NITROUS OXIDE ACETYLENE FLAME.

A 0.5 GRAM SAMPLE WAS REACTED WITH 1.5 ML CONCENTRATED  $\text{HN03}$  AT 90C FOR 30 MINUTES.

AT THIS POINT 0.5 ML CONCENTRATED  $\text{HCL}$  WAS ADDED AND THE DIGESTION WAS CONTINUED AT 90C FOR AN ADDITIONAL 90 MINUTES.

AFTER COOLING, 8 ML OF 1250 PPM  $\text{AL}$  SOLUTION WERE ADDED AND THE SAMPLE SOLUTION WAS DILUTED TO 10 ML BEFORE ASPIRATION.

MERCURY WAS DETERMINED BY THE HATCH AND OTT PROCEDURE WITH SOME MODIFICATIONS. THE METHOD IS DESCRIBED BY JONASSON ET AL. (1973).

A 0.5 GRAM SAMPLE WAS REACTED WITH 20 ML CONCENTRATED  $\text{HNO3}$  AND 1 ML CONCENTRATED  $\text{HCL}$  IN A TEST-TUBE FOR 10 MINUTES AT ROOM TEMPERATURE PRIOR TO 2 HOURS OF DIGESTION WITH MIXING AT 90C IN A HOT WATER BATH.

AFTER DIGESTION, THE SAMPLE SOLUTIONS WERE COOLED AND DILUTED TO 100 ML WITH METAL FREE WATER.

THE  $\text{HG}$  PRESENT WAS REDUCED TO THE ELEMENTAL STATE BY THE ADDITION OF 10 ML  $\text{W/V}$   $\text{SnSO4}$  IN  $\text{M H2SO4}$ .

THE  $\text{HG}$  VAPOUR WAS THEN FLUSHED BY A STREAM OF AIR INTO AN ABSORTION CELL MOUNTED IN THE LIGHT PATH OF AN ATOMIC ABSORTION SPECTROPHOTOMETER.

ABSORPTION MEASUREMENTS WERE MADE AT 253.7 NM.

URANIUM WAS DETERMINED USING A NEUTRON ACTIVATION METHOD WITH DELAYED NEUTRON COUNTING.

WITH THE EXCEPTION OF THE IRRADIATION FACILITY, THE METHOD IS VERY SIMILAR TO THAT USED BY AECL IN PREVIOUS YEARS, A DETAILED DESCRIPTION OF WHICH IS PROVIDED BY BOULANGER ET AL (1975).

A TWO GRAM SAMPLE WAS IRRADIATED FOR 10 SECONDS IN THE TRIGA REACTOR LOCATED AT WASHINGTON STATE UNIVERSITY.

THE OPERATING FLUX WAS  $8 \times 10^{13}$  NEUTRONS/SQUARE CM/SECOND.

AFTER A 10 SECOND DELAY, THE SAMPLE WAS COUNTED FOR 10 SECONDS.

THE COUNTING EQUIPMENT WAS OF AECL DESIGN. CALIBRATION WAS DONE TWICE A DAY OR AS REQUIRED.

ONE STANDARD WAS ANALYSED AFTER EVERY 20 SAMPLES.

TUNGSTEN WAS DETERMINED AS FOLLOWS: A 0.2 GRAM SAMPLE OF STREAM SEDIMENT WAS FUSED WITH 1 GRAM  $\text{KHSO4}$  IN A RIMLESS TEST TUBE AT 575C FOR 15 MINUTES IN A FURNACE. THE COOLED MELT WAS THEN LEACHED WITH 10 ML CONCENTRATED  $\text{HCL}$  IN A WATER BATH HEATED TO 85C. AFTER THE SOLUBLE MATERIAL HAD COMPLETELY DISSOLVED, THE INSOLUBLE MATERIAL WAS ALLOWED TO SETTLE AND AN ALIQUOT OF 5 ML WAS TRANSFERRED TO ANOTHER TEST TUBE. 5 ML OF 20%  $\text{SnCl2}$  SOLUTION WERE THEN ADDED TO THE SAMPLE ALIQUOT, MIXED AND HEATED FOR 10 MINUTES AT 85C IN A HOT WATER BATH. A 1 ML ALIQUOT OF DITHIOL SOLUTION ( 1% DITHIOL IN ISO-AMYL ACETATE) WAS ADDED TO THE TEST SOLUTION AND THE TEST SOLUTION WAS THEN HEATED FFOR 4-6 HOURS AT 80-85C IN A HOT WATER BATH. THE TEST SOLUTION WAS THEN REMOVED FROM THE HOT WATER BATH, COOLED AND 2.5 ML OF KEROSENE ADDED TO DISSOLVE THE GLOBULE SOLUTION WAS MEASURED AT 630 NM USING A SPECTROPHOTOMETER.

FOR THE DETERMINATION OF ANTIMONY A 2-GRAM SAMPLE WAS DIGESTED WITH CONCENTRATED HCL IN A HOT WATER BATH. THE IRON WAS REDUCED TO FE (II) AND THE SB EXTRACTED WITH TRIOCTYL PHOSPHINE OXIDE MIBK AND MEASURED WITH ATOMIC ABSORPTION SPECTROSCOPY WITH BACKGROUND CORRECTION.

FLUORIDE IN LAKE WATER SAMPLES WAS DETERMINED USING AN ORION FLUORIDE ELECTRODE AND A MODEL 404 ORION SPECIFIC ION METER. PRIOR TO MEASUREMENT AN ALIQUOT OF THE SAMPLE WAS MIXED WITH AN EQUAL VOLUME OF TISAB SOLUTION (TOTAL IONIC STRENGTH ADJUSTMENT BUFFER).

HYDROGEN ION ACTIVITY (PH) WAS MEASURED WITH A BROADLEY-JAMES COMBINATION ELECTRODE AND A MODEL 404 ORION SPECIFIC ION METER.

URANIUM IN WATERS WAS DETERMINED BY A LASER-INDUCED FLUOROMETRIC METHOD USING A SCINTREX UA-3 URANIUM ANALYSER. A COMPLEXING AGENT, KNOWN COMMERCIALY AS FLURAN AND COMPOSED OF SODIUM PYROPHOSPHATE AND SODIUM MONOPHOSPHATE, (HALL, G.E.M., 1979) IS ADDED TO PRODUCE THE URANYL PYROPHOSATE SPECIES WHICH FLUORESCES WHEN EXPOSED TO THE LASER. SINCE ORGANIC MATTER IN THE SAMPLE CAN CAUSE UNPREDICABLE BEHAVIOUR, A STANDARD ADDITION METHOD WAS USED. FURTHER, THERE HAVE BEEN INSTANCES AT THE G.S.C. WHERE THE REACTION OF URANIUM WITH FLURAN IS EITHER DELAYED OR SLUGGISH; FOR THIS REASON AN ARBITRARY 24 HOUR TIME DELEAY BETWEEN THE ADDITION OF THE FLURAN AND THE ACTUAL READING WAS INCORPORATED INTO THIS METHOD. IN PRACTICE, 500UL FLURAN SOLUTION WAS ADDED TO A 5ML SAMPLE AND ALLOWED TO STAND FOR 24 HOURS. AT THE END OF THIS PERIOD FLUORESCENCE READINGS WERE MADE WITH THE ADDITION OF 0.0, 0.2 AND 0.4 PPB U. FOR HIGH SAMPLES THE ADDITIONS WERE 0.0, 2.0 AND 4.0 (20UL ALIQUOTS OF 55 OR 550 PPB U WERE USED). ALL READINGS WERE TAKEN AGAINST A SAMPLE BLANK.

THE FOLLOWING TABLE DISPLAYS THE DATA RECORD FORMAT SPECIFICATIONS.  
THE DETECTION LIMITS OF THE ANALYTICAL METHODS ARE ALSO GIVEN WITH  
THE SECOND FIGURE UNDER DETECTION LIMIT USED AS AN ARBITRARILY SET VALUE  
IF THE RESULT FELL BELOW THE DETECTION LIMIT.

FIELD	ELEMENT	CARD	COLUMNS
	MAP	1	01-06
	ID	1	07-12
	UTM ZONE	1	13-14
	UTM EAST (METER)	1	15-20
	UTM NORTH (METER)	1	21-27
	ROCK TYPE	1	28-31
	SAMPLE MATERIAL	1	32
	STREAM WIDTH (DECIMETER)	1	33-35
	STREAM DEPTH (DECIMETER)	1	36-38
	REPLICATE STATUS	1	39-40
	CONTAMINATION	1	41
	BANK TYPE	1	42
	WATER COLOUR	1	43
	FLOW RATE	1	44
	SEDIMENT COLOUR	1	45
	SAMPLE COMPOSITION	1	46-48
	PRECIPITATE IN STREAM	1	49
	DISTINCTIVE PRECIPITATE	1	50
	GENERAL RHYSIOGRAPHY	1	55
	DRAINAGE PATTERN	1	56
	STREAM TYPE	1	57
	STREAM CLASS	1	58
	SOURCE OF WATER	1	59
	AGE	1	70-71

THE ANALYTICAL DATA WERE RECORDED AS FOLLOWS:

ELEMENT	UNITS	CARD	COLUMNS	DETECTION LIMIT
SEDIMENT				
ZN	PPM	2	21-25	2 1
CU	PPM	2	26-30	2 1
PB	PPM	2	31-35	2 1
NI	PPM	2	36-40	2 1
CO	PPM	2	41-45	2 1
AG	PPM	2	46-50	0.2 0.1
MN	PPM	2	51-55	5 2
AS	PPM	2	56-60	1.0 0.5
MO	PPM	2	61-65	2 1
FE	PCT	2	66-70	0.02 0.01
HG	PPB	2	71-75	10 5
U	PPM	3	21-25	0.2 0.1
W	PPM	3	26-30	2 1
SB	PPM	3	36-40	0.2 0.1
F-W	PPB	4	26-30	20 10
PH		4	31-35	
U-W	PPB	4	36-40	0.05 0.002

REFERENCES

BOULANGER, A., EVANS, D.J.R. AND RABY, B.F. (1975) URANIUM ANALYSIS BY NEUTRON ACTIVATION DELAYED NEUTRON COUNTING: PROC. OF THE 7TH ANNUAL SYMP. OF CANADIAN MINERAL ANALYSTS. THUNDER BAY, ONTARIO, SEPT. 22-23, 1975.

FICKLIN, W.H. (1970) A RAPID METHOD FOR THE DETERMINATION OF FLUORIDE IN ROCKS AND SOILS, USING AN ION SELECTIVE ELECTRODE. U.S. GEOL. SURV. PAPER 700C PP. C186-188.

GARRETT, R.G. (1974) FIELD DATA ACQUISITION METHODS FOR APPLIED GEOCHEMICAL SURVEYS AT THE GEOLOGICAL SURVEY OF CANADA: GEOL SURV. CAN. PAPER 74-52.

ASLIN, G.E.M. (1976) THE DETERMINATION OF ARSENIC AND ANTIMONY IN GEOLOGICAL MATERIALS BY FLAMELESS ATOMIC ABSORPTION SPECTROPHOTOMETER JOURNAL OF GEOCHEMICAL EXPLORATION, VOL. 6, PP. 321-330.

HALL, G.E.M. (1979) A STUDY OF THE STABILITY OF URANIUM IN WATERS COLLECTED FROM VARIOUS GEOLOGICAL ENVIRONMENTS IN CANADA; IN CURRENT RESEARCH, PART A, GEOL. SURV. CAN. PAPER 79-1A, P. 361-365.

JONASSON, I.R., LYNCH, J.J. AND TRIP, L.J. (1973) FIELD AND LABORATORY METHODS USED BY THE GEOLOGICAL SURVEY OF CANADA IN GEOCHEMICAL SURVEYS: NO. 12, MERCURY IN ORES, ROCKS, SOILS, SEDIMENTS AND WATER: GEOL. SURV. CAN. PAPER 73-21.

DATA LIST LEGEND

MAP- NATIONAL TOPOGRAPHIC SYSTEM(NTS)- LETTERED SIXTEENTH  
(SCALE 1:500000). PART OF SAMPLE NUMBER

SAMPLE- REMAINDER OF SAMPLE NUMBER- YEAR(2), FIELD CREW(1),  
SAMPLE SEQUENCE NUMBER(3)

UTM COORDINATES- UNIVERSAL TRANSVERSE MERCATOR(UTM) COORDINATE  
SYSTEM- SAMPLE COORDINATES

ZN- ZONE

EAST- EASTINGS (METERS)

NORTH- NORTHINGS (METERS)

ROCK TYPE- MAJOR ROCK TYPE OF CATCHMENT AREA

AGE- STRATIGRAPHIC AGE OF ROCK TYPE

WD- WIDTH OF STREAM (DECIMETER) AT NEAREST SAMPLE SITE

DT- DEPTH OF STREAM SAMPLED TO NEAREST DECIMETER

SAMP- TYPE OF MATERIAL SAMPLED

RP ST- REPLICATE STATUS- RELATIONSHIP OF SAMPLE WITH RESPECT  
TO OTHERS WITHIN THE SURVEY

CONT- CONTAMINATION

BANK- BANK TYPE

WCOL- WATER COLOUR AND SUSPENDED LOAD

RATE- WATER FLOW RATE

SCOL- PREDOMINANT SEDIMENT COLOUR

SMP CMP- SAMPLE COMPOSITION- BULK MECHANICAL COMPOSITION OF  
SAND, FINES, ORGANICS RESPECTIVELY

PPPS- PRECIPITATE OR STAIN ON SEDIMENTS AT SAMPLE SITE

PRPB- DISTINCTIVE PRECIPITATE, STAIN, WEATHERING, BLOOMS ON  
ROCKS IN THE IMMEDIATE CATCHMENT AREA

PHYS- GENERAL PHSYIOGRAPHY

PATT- DRAINAGE PATTERN

TYPE- STREAM TYPE

CLSE- STREAM CLASS

SRCE- SOURCE OF WATER



ROCK TYPE :	(TILL 44) - TILL, GRAVEL, SAND, SILT, ALLUVIUM	AGE :	04- PROTEROZOIC (600-2400 M.Y.) 10- PALEOZOIC UNDIVIDED 17- SILURIAN-DEVONIAN 23- PENNSYLVANIAN - PERMAIN 32- TRIASSIC 34- JURASSIC 36- CRETACEOUS 41- MESOZOIC - CENOZOIC 44- QUATERNARY 50- UNKNOWN
	(RYLT 41) - OOTSA LAKE GROUP : RHYOLITE, DACITE, TRACHYTE, SANDSTONE, SHALE, CONGLOMERATE		
	(CGLM 41) - SUSTUT GROUP: CONGLOMERATE, SHALE, GREYWACKE		
	(SHLE 36) - SKEENA GROUP, RED ROSE FORMATION : SHALE, GREYWACKE, CONGLOMERATE, COAL, VOLCANIC BRECCIA		
	(BSLT 34) - TELKWA AND NILKITKWA FORMATIONS : BASALT, ANDESITE, BRECCIA, TUFF, SHALE, SILTSTONE	SAMP :	1- STREAM BED SEDIMENT 6- SIMULTANEOUS STREAM WATER AND SEDIMENT
	(ANDS 32) - TAKLA GROUP : ANDESITE, BASALT, TUFF, BRECCIA, CONGLOMERATE, GREYWACKE, SHALE, LIMESTONE	RP ST :	00- ROUTINE REGIONAL SAMPLE 10- FIRST OF FIELD DUPLICATE 20- SECOND OF FIELD DUPLICATE
	(LMSN 23) - CACHE CREEK GROUP : RIBBON CHERT, BLACK ARGILLITE, LIMESTONE, GREENSTONE		
	(LMDM 17) - LIMESTONE, DOLOMITE, SANDY DOLOMITE, QUARTZITE, SHAL	CONT :	0- NONE 1- POSSIBLE 2- PROBABLE 3- DEFINITE
	(GRNS 10) - GREENSTONE, ANDESITIC VOLCANIC ROCKS, ARGILLITE, SHALE, LIMESTONE		
	(MSDM 1) - UNDIVIDED METASEDIMENTARY AND SEDIMENTARY ROCKS OF HADRYNIAN TO LOWER DEVONIAN AGE	BANK :	0- UNDEFINED UNCONSOLIDATED MATERIAL 1- ALLUVIAL 2- COLLUVIAL 3- GLACIAL TILL, TILLITE 4- GLACIAL OUTWASH, MORaine 6- TALUS, SCREE 7- ORGANIC PREDOMINANT
	(PLLT 04) - INGENIKA GROUP : UNDIVIDED PHYLLITE, SCHIST, GRIT, LIMESTONE		
	(GRNG 50) - WOLVERINE METAMORPHIC COMPLEX : GRANITOID GNEISS, PEGMATITE, SCHIST, AMPHIBOLITE, QUARTZITE	WCOL :	0- CLEAR 1- BROWN TRANSPARENT 3- BROWN CLOUDY
	(GRNT 41) - NAYER INTRUSIONS, TOPLEY INTRUSIONS, DUCKLING CREEK SYENITE COMPLEX, HOGEM BATHOLITH, OMINECA INTRUSIONS, AND SIMILAR GRANITIC ROCKS: QUARTZ DIORITE, DIORITE, QUARTZ MONZONITE, GRANODIORITE, AND SYENITE WITH MINOR GRANITE, PEGMATITE, AND APLITE	RATE :	0- ZERO 1- SLOW 2- MODERATE 3- FAST 4- TORRENTIAL
	(SRPM 41) - TREMBLEUR INTRUSIONS AND SIMILAR ULTRAMAFIC BODIES: PERIDOTITE, DUNITE, PYROXENITE, AND SERPENTINITE	SCOL :	0- UNKNOWN 1- RED, BROWN 2- WHITE, BUFF 3- BLACK 5- GREEN 6- GREY, BLUE-GREY

DATA LEGEND CONT.

SMP CMP :       0- ABSENT  
                   1- MINOR <33%  
                   2- MEDIUM <33-67%  
                   3- MAJOR > 67%

PPPS :           0- NONE  
                   1- RED, BROWN

PRPB :           0- FEATURELESS  
                   1- RED, BROWN

PHYS :           1- MUSKEG, SWAMPLAND  
                   2- PENEPLAIN, PLATEAU  
                   3- HILLY, UNDULATING  
                   4- MOUNTAINOUS MATURE  
                   5- MOUNTINOUS YOUTHFUL

PATT :           0- POORLY DEFINED, HAPHAZARD  
                   1- DENDRITIC  
                   2- HERRING BONE  
                   3- TRELLISED (FOLDED)

TYPE :           1- PERMANENT, CONTINUOUS  
                   2- INTERMITTENT, SEASONAL  
                   3- RE-EMERGENT, DISCONTINUOUS  
                   4- UNKNOWN

CLSE :           3- TERTIARY  
                   4- QUARERNARY

SRCE :           1- GROUNDWATER  
                   2- SNOW MELT OR SPRING RUN-OFF

ZN- ZINC BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 CU- COPPER BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 PB- LEAD BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 NI- NICKEL BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 CO- COBALT BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 AG- SILVER BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 MN- MANGANESE BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 AS- ARSENIC BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 MO- MOLYBDENUM BY ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 FE- IRON BY ATOMIC ABSORPTION SPECTROSCOPY (%)  
 HG- MERCURY BY ATOMIC ABSORPTION SPECTROSCOPY (PPB)  
 U- URANIUM BY DELAYED NEUTRON ACTIVATION (PPM)  
 W- TUNGSTON BY COLORIMETRY USING DITHIOL (PPM)  
 SB- ANTIMONY MIBK SOLVENT EXTRACTION ATOMIC ABSORPTION SPECTROSCOPY (PPM)  
 F-W- FLOURINE IN WATERS BY FISSION TRACK (PPB)  
 PH- PH BY COMBINATION GLASS - CALOMEL ELECTRODE  
 U-W- URANIUM IN WATERS BY SCINTREX (PPB)