



DESCRIPTIVE NOTES

**Geological**

Almost all of the map-area is underlain by Palaeozoic rocks that range in age from Early Ordovician to Late Devonian. Volcanic rocks in the extreme southeastern part of the map-area are tentatively correlated with the Charlotte Group rocks of probable Proterozoic age. The northwestern third of the map-area is underlain by a widespread sequence of undifferentiated Lower Palaeozoic light to dark slates, argillites and impure quartzites of the Charlotte Group.

In the Oak Bay region, the base of the Silurian is marked by a belt of conglomerates that grades upward into an assemblage of greywackes, quartzite, siltstone and argillite. Andesite, rhyolite, diatite and basalt flows, breccias and tuffs predominate in the upper part of the Silurian section between Bocabec Bay and Lake Utopia. Volcanic and sedimentary units on the Mascarene-Lettie Peninsula are highly deformed and intensely sheared, in contrast to generally less deformed unmetamorphosed rocks of similar lithology in the area northwest of the Magaguadavic River.

Eastport-type sedimentary rocks of possible Lower Devonian age are intruded by rhyolite and andesite volcanic necks near Chamcook Lake. The Upper Devonian Perry Formation consists of extensive terrestrial deposits of red conglomerates and sandstones, and minor interbedded basalt flows. Conglomerates containing granitic boulders and clasts of Silurian and pre-Silurian rocks are in most cases undeformed and rest unconformably on older rocks or in fault contact with them.

Lower Palaeozoic volcanic and sedimentary rocks are intruded by Ordovician gabbro-norite bodies, and Middle Devonian granitic batholiths and stocks. The large St. George batholith underlies the northeastern part of the map-area and is composed mainly of granite, quartz monzonite and adamellite. Three smaller stocks of similar composition intrude the Lower Palaeozoic sedimentary rocks of the Charlotte Group. Granitic rocks immediately east of Oak Bay are intimately associated with gabbro, diorite and diabase that are similar to, but may significantly post-date the gabbro intrusives at St. Stephen, Silurian and Lower Devonian sedimentary and volcanic rocks are locally intruded by gabbro and diabase dykes, sills and stocks.

The region has been heavily glaciated. Eskers, glacial striae, and stoss and lee slopes indicate a general southeasterly direction of ice movement. Double sets of striae in the area south of St. George reflect at least two ice advances. An older set bearing 145 degrees is combined with a younger set that averages 110 degrees. Thick deposits of glacial till and fluvio-glacial sands and gravels cover most of the area and have markedly altered the pre-Pleistocene drainage pattern on the lower parts of the Magaguadavic River.

The principal mineral deposits in the district are massive, vein and disseminated copper-nickel deposits associated with the gabbroic intrusions, and copper-lead-zinc disseminated and vein-type deposits in the Silurian volcanic rocks. Molybdenum, tin and gold occurrences appear to be associated with some of the granitic bodies.

Map 1 should be consulted for further details on the geology and economic geology of the district.

**Geochemical**

The data recorded on this map are based on the analyses of 1,056 samples of fine-grained sediment collected from the channels of rivers and streams, and from rivulets flowing from springs and seeps. Where possible the active channels were sampled, but in a few cases bank material or residual sediment of dried-up streams was used. Sediment samples from poorly drained areas of muskeg and beaver workings contained abundant decomposed organic matter. All samples were collected during the 1966 field season. An attempt was made to maintain a uniform sample density, but this was frequently not achieved because of irregularities in the drainage network and the absence of an actively deposited silt-limed fraction in streams draining areas underlain by coarse glacial till. All streams and rivers were traversed on foot, and samples were collected, where possible, at intervals of 1,000 to 1,500 feet.

Sediment samples were air dried prior to shipment to Bondar-Clegg & Company Limited, Ottawa, Ontario for laboratory analysis. Zinc was determined by atomic absorption analysis after extraction from the -80 mesh sieve fraction by hot HNO<sub>3</sub> - HCl digestion. Values shown on this map are expressed in parts per million. The subdivisions used on the map are arbitrary, and the lowest subdivision can be taken to represent the regional background.

The zinc content of the stream sediments ranges from 12 ppm to 5,300 ppm. The frequency distribution of zinc shows that 52.5 per cent of the analyses exceed the normal background value of 70 ppm, and 11.9 per cent of the analyses exceed 120 ppm. In such an area of widely diverse rock types, important variations in zinc background may be related to local variations in bedrock lithology, proximity to disseminated deposits or sparsely mineralized rocks, or to concentrations of strongly adsorbing organic matter and/or hydrous manganese-iron bodies. Manganeseiferous stream sediments are commonly enriched in zinc. The strong positive correlation between the total manganese and total zinc contents of 1,025 stream sediments from unmineralized terrain is shown in Figure 2. The calculation and use of a single zinc threshold value will produce a number of manganese dependent anomalies that are unrelated to mineral deposits. Samples from streams draining mineralized areas generally have Zn/Mn ratios higher than predicted by the data shown in Figure 2.

Copper-lead-zinc occurrences on the Mascarene-Lettie Peninsula are marked by higher than normal contents of Zn in nearby streams. A number of isolated zinc anomalies occur throughout the district in previously unexplored terrain, and these should receive further investigation. Sediments from streams draining the coastal region between the Digdeguash and Magaguadavic Rivers have produced anomalous zinc values (200 ppm to 1,300 ppm) that are unrelated to high manganese contents, known deposits or contaminating agencies. These streams drain an area that is underlain by andesite, rhyolite and basalt flows and intercalated red shales and siltstones of Silurian age.

The zinc contents of stream and spring sediments shown on this map should be interpreted with respect to the geological environment (Map 1) and compared with the contents of identical elements in stream sediments on Maps 2 to 5 inclusive.

**Figure 1** Histogram of zinc concentrations in 1056 samples of stream sediment.

**Figure 2** Correlation between total zinc and total manganese contents of 1025 stream sediments from unmineralized terrain.

**Metal and Mineral Symbols**

Arsenic ..... As	Silver ..... Ag	Hematite ..... hem
Bismuth ..... Bi	Zinc ..... Zn	Malachite ..... mal
Copper ..... Cu	Arsenopyrite ..... asp	Pyrrhotite ..... pyr
Gold ..... Au	Chalcopyrite ..... cp	Pyrite ..... py
Lead ..... Pb	Fluorite ..... fl	Quartz ..... qtz
Molybdenum ..... Mo	Galenite ..... gn	Sphalerite ..... sp
Nickel ..... Ni		

**INDEX TO MINING PROPERTIES AND PROSPECTS**

1. Atlantic Nickel (Rogers Farm): cp, po, pm.	9. Rollingdam Molybdenite: Mo, fl.
2. Clark Farm (A' Zone): cp, po.	10. Mt. Blair: cp, fl, mal, hem.
3. Hall Carroll: po, cp.	11. Oliver - Cameron: qtz, cp, Bi, Au, Ag, mal.
4. Dennis Stream (W' Zone): cp, po.	12. T. Dick: qtz, py, cp, po, gn.
5. Grant Farm: cp, po.	13. Lettie: qtz, cp, gn.
6. 'C' Zone: cp, po.	14. Simpson: cp, gn, py.
7. Union Bridge - St. Croix River: cp, po.	15. Daniel Hart Farm: qtz, cp, po, gn.
8. Moores Mills - qtz, asp, gn, sp.	

Field Work by W.J. Wolfe, M. Mason, G.J. Mazerolle, J.M. Robbins, R.G. Garnett and D.K.J. MacKinnon

Analyses by Bondar-Clegg & Company Limited, Ottawa, Ontario.

**MAP 3**  
REPORT OF INVESTIGATIONS NO 4  
ZINC CONTENT OF STREAM AND SPRING SEDIMENTS  
SOUTHWESTERN CHARLOTTE COUNTY, NEW BRUNSWICK

SCALE 1:100,000  
1.25 inches to 2 miles approximately

Location of known mineral occurrences (Symbols indicate principal metals)

Mining properties and prospects (see index)

**Published, 1967**  
Copies of this map may be obtained from the Mines Division, Department of Natural Resources, Fredericton, New Brunswick