

NEW BRUNSWICK
MINES DIVISION
DEPARTMENT OF NATURAL RESOURCES

DESCRIPTIVE NOTES

Geological

Almost all of the map-area is underlain by Paleozoic rocks that range in age from Early Ordovician to Late Devonian. Volcanic rocks in the extreme southeastern part of the map-area are tentatively correlated with Collibrook Group rocks of probable Proterozoic age. The northwestern third of the map-area is underlain by a widespread sequence of undifferentiated Lower Paleozoic light to dark slates, argillites and impure quartzites of the Charlotte Group.

In the Oak Bay region, the base of the Silurian is marked by a belt of conglomerate that grades upward into an assemblage of greywacke, quartzite, siltstone and argillite. Andesite, rhyolite, dacite and basalt flows, breccias and tuffs predominate in the upper part of the Silurian section between Bocabec Bay and Lake Utopia. Volcanic and sedimentary units on the Macareno-Lafite Peninsula are highly deformed and intensely sheared, in contrast to generally less deformed unmetamorphosed rocks of similar lithology in the area northwest of the Magaguadavic River.

Eastport-type sedimentary rocks of possible Lower Devonian age are intruded by rhyolite and andesite volcanic necks near Chamoock Lake. The Upper Devonian Perry Formation consists of extensive terrarial deposits of red conglomerate and sandstone, and minor intercalated basalt flows. Conglomerates containing granitic boulders and clasts of Silurian and pre-Silurian rocks are in most cases undeformed and rest unconformably on older rocks or in fault contact with them.

Lower Paleozoic volcanic and sedimentary units are intruded by Ordovician gabbro-norite bodies, and Middle Devonian granitic batholiths and stocks. The large St. George batholith underlies the northeastern part of the map-area and is composed mainly of granite, quartz monzonite and adamellite. Three smaller stocks of similar composition intrude the Lower Paleozoic sedimentary rocks of the Charlotte Group. Granitic rocks immediately east of Oak Bay are intimately associated with gabbro, diorite and diabase that are similar to, but may significantly post-date the gabbro intrusives at St. Stephen. Silurian and Lower Devonian sedimentary and volcanic rocks are locally intruded by gabbro and diabase dykes, sills and stocks.

The region has been heavily glaciated. Eskers, glacial striae, and steep and lee slopes indicate a general southeasterly direction of ice movement. Double sets of striae in the area south of St. George reflect at least two ice advances. An older set bearing 145 degrees is combined with a younger set that averages 110 degrees. Thick deposits of glacial till and fluvio-glacial sands and gravels cover most of the area and have markedly altered the pre-Pleistocene drainage pattern on the lower parts of the Magaguadavic River.

The principal mineral deposits in the district are massive, vein and disseminated copper-nickel deposits associated with the gabbroic intrusions, and copper-lead-zinc disseminated and vein-type deposits in the Silurian volcanic rocks. Molybdenum, tin and gold occurrences appear to be associated with some of the granitic bodies.

Map 1 should be consulted for further details on the geology and economic geology of the district.

Geochemical

The data recorded on this map are based on the analyses of 1,056 samples of fine-grained sediment collected from the channels of rivers and streams, and from rivulets flowing from springs and seeps. Where possible the active channels were sampled, but in a few cases bank material or residual sediment of dried-up streams was used. Sediment samples from poorly drained areas of muskeg and field season. An attempt was made to maintain a uniform sample density, but this was frequently not achieved because of irregularities in the drainage network and the absence of an actively deposited alluvial fraction in streams draining areas underlain by coarse glacial till. All streams and rivers were traversed on foot, and samples were collected, where possible, at intervals of 1,000 to 1,500 feet.

Sediment samples were air dried prior to shipment to Bondar-Clegg & Company Limited, Ottawa, Ontario for laboratory analysis. The 40 mesh sieve fraction was used for the colorimetric determination of molybdenum after extraction by pyrosulphate fusion and HCl fusion. Values shown on this map are expressed in parts per million. The subdivisions used on the map are arbitrary, and the lowest subdivision can be taken to represent the regional background.

The molybdenum content of the stream sediments ranges from non-detectable (less than 1 ppm) to 40 ppm. The background for the whole district is probably 1 ppm or less, but important background variations are related to local variations in bedrock lithology, and the normal molybdenum content of stream sediments in areas underlain by granitic rocks may be as high as 4 to 5 ppm. The frequency distribution of molybdenum data shows that 44.0 per cent of the analyses exceed the detection limit of 1 ppm, and 12.5 per cent of the analyses exceed 5 ppm.

Many of the streams draining areas underlain by granitic stocks and batholiths and their associated metamorphic aureoles contain above normal amounts of molybdenum in the stream sediments. Anomalous amounts of molybdenum are spatially related to the margins of small granitic stocks west of the St. George batholith in the area directly north of the Bocabec River. Other streams of interest are those draining the general area underlain by granitic and contact rocks along the southern margin of the St. George batholith between the Digdeguash River and Lake Utopia. Anomalous Mo, Cu, Zn values may reflect mineralization associated with the south contact of the Clarence stock to the northeast of the Digdeguash River.


There is no general correlation of the molybdenum and manganese contents of the sediments. Although the presence of large concentrations of hydrous manganese-iron oxides may be a factor in the localization of molybdenum in some of the anomalous streams, many manganese-rich sediments have molybdenum values of background or lower. Low molybdenum environments apparently place a limiting restriction on the possible amount of Mo accumulation in manganese oxides.

The molybdenum contents of stream and spring sediments shown on this map should be interpreted with respect to the geological environment (Map 1) and compared with the contents of individual elements in stream sediments on Maps 2 to 8 inclusive.

INDEX TO MINING PROPERTIES AND PROSPECTS

1. Atlantic Nickel (Rodgers Farm): cp, po, pn.	9. Rollington Molybdenite: mo, fl.
2. Clark Farm ('A' Zone): cp, po.	10. Mt. Blair: cp, fl, mal, hem.
3. Hall Carroll: po, cp.	11. Oliver - Cameron: qts, cp, Bi, Au, Ag, mal.
4. Dennis Stream ('N' Zone): cp, po.	12. T. Dick: qts, py, cp, po, gn.
5. Great Farm: cp, po.	13. Lefter: qts, cp, gn.
6. 'C' Zone: cp, po.	14. Simpson: cp, gn, py.
7. Union Bridge - St. Croix River: cp, po.	15. Daniel Hall Farm: qts, cp, po, gn.
8. Moore's Mill: qts, asp, gn, sp.	

Field work by W.J. Wolfe, M. Mason, G.J. Mazerolle, J.M. Robbins, R.G. Garnett and D.K.J. MacKinnon
Analyses by Bondar-Clegg and Company Limited, Ottawa, Ontario.



Published, 1967
Copies of this map may be obtained from the
Mines Division, Department of Natural Resources,
Fredericton, New Brunswick