

DESCRIPTIVE NOTES

Geological

Almost all of the map-area is underlain by Palaeozoic rocks that range in age from Early Ordovician to Late Devonian. Volcanic rocks in the extreme southeastern part of the map-area are tentatively correlated with the Charlotte Group rocks of probable Proterozoic age. The northwestern third of the map-area is underlain by a widespread sequence of undifferentiated Lower Palaeozoic light to dark slates, argillites and impure quartzites of the Charlotte Group.

In the Oak Bay region, the base of the Silurian is marked by a belt of conglomerate that grades upward into an assemblage of greywacke, quartzite, siltstone and argillite. Andesite, rhyolite, dacite and basalt flows, breccias and tuffs predominate in the upper part of the Silurian section between Bocabe Bay and Lake Utopia. Volcanic and sedimentary units on the Macarone-Little Peninsula are highly deformed and intensely sheared, in contrast to generally less deformed unmetamorphosed rocks of similar lithology in the area northwest of the Magaguadavic River.

Eastport-type sedimentary rocks of possible Lower Devonian age are intruded by rhyolite and andesite volcanic necks near Chamcook Lake. The Upper Devonian Perry Formation consists of extensive terrestrial deposits of red conglomerate and sandstone, and minor intercalated basalt flows. Conglomerates containing granitic boulders and clasts of Silurian and pre-Silurian rocks are in most cases undeformed and rest unconformably on older rocks or in fault contact with them.

Lower Palaeozoic volcanic and sedimentary units are intruded by Ordovician gabbro-norite bodies, and Middle Devonian granitic batholiths and stocks. The large St. George batholith underlies the northeastern part of the map-area and is composed mainly of granite, quartz monzonite and adamellite. Three smaller stocks of similar composition intrude the Lower Palaeozoic sedimentary rocks of the Charlotte Group. Granitic rocks immediately east of Oak Bay are intimately associated with gabbro, diorite and diabase that are similar to, but may significantly post-date the gabbro intrusives at St. Stephen. Silurian and Lower Devonian sedimentary and volcanic rocks are locally intruded by gabbro and diabase dykes, sills and stocks.

The region has been heavily glaciated. Eskers, glacial striae, and stoss and lee slopes indicate a general southeasterly direction of ice movement. Double sets of striae in the area south of St. George reflect at least two ice advances. An older set bearing 145 degrees is combined with a younger set that averages 110 degrees. Thick deposits of glacial till and fluvio-glacial sands and gravels cover most of the area and have markedly altered the pre-Pleistocene drainage pattern on the lower parts of the Magaguadavic River.

The principal mineral deposits in the district are massive, vein and disseminated copper-nickel deposits associated with the gabbroic intrusions, and copper-lead-zinc disseminated and vein-type deposits in the Silurian volcanic rocks. Molybdenum, tin and gold occurrences appear to be associated with some of the granitic bodies.

Map 1 should be consulted for further details on the geology and economic geology of the district.

Geochemical

The data recorded on this map are based on the analyses of 1,056 samples of fine-grained sediment collected from the channels of rivers and streams, and from rivulets flowing from springs and seeps. Where possible the active channels were sampled, but in a few cases bank material or residual seeps. Sediment samples from poorly drained areas of muskeg and heavier workings contained abundant decomposed organic matter. All samples were collected during the 1966 heavy working season. An attempt was made to maintain a uniform sample density, but this was frequently not achieved because of irregularities in the drainage network and the absence of an actively deposited silt-sized fraction in streams draining areas underlain by coarse glacial till. All streams and rivers were traversed on foot, and samples were collected, where possible, at intervals of 1,000 to 1,500 feet.

Sediment samples were air dried prior to shipment to Bondar-Clegg & Company Limited, Ottawa, Ontario for laboratory analysis. Manganese was determined by atomic absorption analysis after extraction from the -80 mesh sieve fraction by hot HNO₃ - HCl digestion. Values shown on this map are expressed in parts per million. The subdivisions used on the map are arbitrary, and the lowest subdivision can be taken to represent the regional background.

The manganese content of the stream sediments ranges from 20 ppm to 10,000 ppm. The frequency distribution of manganese shows that 51.8 per cent of the analyses exceed the normal background value of 800 ppm, and 13.7 per cent of the analyses exceed 2,000 ppm. Local variations in stream gradient, and/or oxidation potential may lead to erratic distribution of manganese in the sediments. The average pH and/or oxidation potential may vary between 6.00 ppm and 1,000 ppm according to the type of underlying bedrock lithology. Isolated samples that are greatly enriched in manganese may reflect the existence of nearby manganiferous sedimentary rocks.

Several heavy metals, including Cu, Pb, Zn and Ni are strongly adsorbed and coprecipitated by manganese hydroxides and oxides. Zinc concentration is most markedly affected by the presence of manganese (see Figure 2), and this feature may lead to false anomalies or abnormal enhancement of valid anomalies.

The manganese contents of stream and spring sediments shown on this map should be interpreted with respect to the geological environment (Map 1) and compared with the contents of individual elements in stream sediments on Maps 2 to 8 inclusive.


Metal and Mineral Symbols

Arsenic	As	Silver	Ag	Hematite	hem
Bismuth	Bi	Malachite	mal	Malachite	mal
Copper	Cu	Pentlandite	pn	Pentlandite	pn
Gold	Au	Arsenopyrite	asp	Pyrite	py
Lead	Pb	Chalcopyrite	cp	Pyrrhotite	pyr
Molybdenum	Mo	Fluorite	fl	Quartz	qtz
Nickel	Ni	Galena	gn	Sphalerite	sp

INDEX TO MINING PROPERTIES AND PROSPECTS

- Atlantic Nickel (Rodgers Farm): cp, po, pn.
- Clark Farm (A' Zone): cp, po.
- Hall Carroll: po.
- Dennis Stream (N' Zone): cp, po.
- Grant Farm: cp, po.
- 'C' Zone: cp, po.
- Union Bridge - St. Croix River: cp, po.
- Moore Mills - qtz, asp, gn.
- Rollingdam Molybdenite: Mo, fl.
- Mr. Blair: cp, fl, mal, hem.
- Oliver - Cameron: qtz, cp, Bi, Au, Ag, mal.
- T. Dick: qtz, py, qtz, po, gn.
- Leiter: qtz, cp, gn.
- Simpson: sp, gn, py.
- Daniel Hart Farm: qtz, cp, po, gn.

Field work by W.J. Wolfe, M. Mason, G.J. Mazerolle, J.M. Robbins, R.G. Garnett and D.K.J. MacKinnon.
Analyses by Bondar-Clegg & Company, Limited, Ottawa, Ontario.



Published, 1967
Copies of this map may be obtained from the
Mines Division, Department of Natural Resources,
Fredericton, New Brunswick




Figure 1. Histogram of manganese concentrations in 1056 samples of stream sediment.




Figure 2. Correlation between total zinc and total manganese contents of 1025 stream sediments from unmineralized terrain.

MAP 5
REPORT OF INVESTIGATIONS NO. 4
MANGANESE CONTENT OF STREAM AND SPRING SEDIMENTS
SOUTHWESTERN CHARLOTTE COUNTY, NEW BRUNSWICK

Sample location
● 1250 Concentration, 900 to 1,999 parts per million
● 3000 Concentration, 2,000 parts or greater per million

SCALE 1:100,000
1.25 inches to 2 miles approximately

Location of known mineral occurrences
(Symbols indicate principal metals)

Mining properties and prospects (see index)