

NEW BRUNSWICK
MINES DIVISION
DEPARTMENT OF NATURAL RESOURCES

DESCRIPTIVE NOTES

Geological

Almost all of the map-area is underlain by Paleozoic rocks that range in age from Early Ordovician to Late Devonian. Volcanic rocks in the extreme southeastern part of the map-area are tentatively correlated with the Charlotte Group rocks of probable Proterozoic age. The northwestern third of the map-area is underlain by a widespread sequence of undifferentiated Lower Paleozoic light to dark slates, argillites and impure quartzites of the Charlotte Group.

In the Oak Bay region, the base of the Silurian is marked by a belt of conglomerate that grades upward into an assemblage of greywacke, quartzite, siltstone and argillite. Andesite, rhyolite, dacite and basalt flows, breccias and tuffs predominate in the upper part of the Silurian section between Bonabec Bay and Lake Utopia. Volcanic and sedimentary units on the Mascaene-Lettie Peninsula are highly deformed and intensely sheared, in contrast to generally less deformed unmetamorphosed rocks of similar lithology in the area northwest of the Magaguadavic River.

Eastport-type sedimentary rocks of possible Lower Devonian age are intruded by rhyolite and andesite volcanic necks near Chamcook Lake. The Upper Devonian Perry Formation consists of extensive terrestrial deposits of red conglomerate and sandstone, and minor intercalated basalt flows. Conglomerates containing granitic boulders and clasts of Silurian and pre-Silurian rocks are in most cases undeformed and rest unconformably on older rocks or in fault contact with them.

Lower Paleozoic volcanic and sedimentary units are intruded by Ordovician gabbro-norite bodies, and Middle Devonian granitic batholiths and stocks. The large St. George batholith underlies the northeastern part of the map-area and is composed mainly of granite, quartz monzonite and adamellite. The Three smaller stocks of similar composition intrude the Lower Paleozoic sedimentary rocks of the Charlotte Group. Granitic rocks immediately east of Oak Bay are intimately associated with gabbro, diorite and diabase that are similar to, but may significantly post-date the gabbro intrusives at St. Stephen. Silurian and Lower Devonian sedimentary and volcanic rocks are locally intruded by gabbro and diabase dykes, sills and stocks.

The region has been heavily glaciated. Eskers, glacial striae, and stoss and lee slopes indicate a general southeasterly direction of ice movement. Double sets of striae in the area south of St. George reflect at least two ice advances. An older set bearing 145 degrees is combined with a younger set that averages 110 degrees. Thick deposits of glacial till and fluvio-glacial sands and gravels cover most of the area and have markedly altered the pre-Pleistocene drainage patterns on the lower parts of the Magaguadavic River.

The principal mineral deposits in the district are massive, vein and disseminated copper-nickel deposits associated with the gabbro intrusions, and copper-lead-zinc disseminated and vein-type deposits in the Silurian volcanic rocks. Molybdenum, tin and gold occurrences appear to be associated with some of the granitic bodies.

Map 1 should be consulted for further details on the geology and economic geology of the district.

Geochemical

The data recorded on this map are based on the analyses of 546 samples of fine-grained sediment collected from the channels of rivers and streams, and from rivulets flowing from springs and seeps. Where possible the active channels were sampled, but in a few cases bank material or residual sediment of dried-up streams was used. Sediment samples from poorly drained areas of muskeg and heaver workings contained abundant decomposed organic matter. All samples were collected during the 1966 field season. An attempt was made to maintain a uniform sample density, but this was frequently not achieved because of irregularities in the drainage network and the absence of an actively deposited silt-sized fraction in streams draining areas underlain by coarse glacial till. All streams and rivers were traversed on foot, and samples were collected, where possible, at intervals of 1,000 to 1,500 feet.

Sediment samples were air dried prior to shipment to Bondar-Clegg & Company Limited, Ottawa, Ontario for laboratory analysis. The arsenic content of the -80 mesh sieve fraction was determined by a modified Gutzeit method using a potassium hydroxide fusion. Values shown on this map are expressed in parts per million. The subdivisions used on the map are arbitrary, and the lowest subdivision can be taken to represent the regional background.

The arsenic content of the stream sediments ranges from non-detectable (less than 3 ppm) to 220 ppm. The frequency distribution of arsenic data shows that 52.2 per cent of all analyses exceed the normal background value of 5 ppm, and 12.4 per cent of the analyses exceed 30 ppm. In such an area of widely diverse rock types, important variations in arsenic background may be related to local variations in bedrock lithology, proximity to sparsely mineralized arsenopyrite-bearing rocks, or to concentrations of strongly adsorbing organic matter and/or hydrous manganese-iron oxides.

Higher than average contents of arsenic in stream sediments are commonly related to areas underlain by granitic stocks and batholiths and their associated metamorphic aureoles. In this respect, the overall dispersion pattern of arsenic is similar to that of molybdenum. Anomalous amounts of arsenic are spatially associated with small granitic stocks and mixed granitic and dioritic rocks west of the St. George batholith in the area directly north of the Bonabec River. Isolated arsenic high in sediments from Gallop Stream and the Waweg River system may reflect the presence of arsenopyrite-bearing quartz veins in sediments of the Charlotte Group.

The arsenic contents of stream and spring sediments shown on this map should be interpreted with respect to the geological environment (Map 1) and compared with the contents of individual elements in stream sediments on Maps 2 to 8 inclusive.


Metal and Mineral Symbols

Arsenic..... As	Silver..... Ag	Hematite..... hem
Bismuth..... Bi	Zinc..... Zn	Malachite..... mal
Copper..... Cu	Antimony..... Sb	Pentlandite..... pn
Gold..... Au	Arsenopyrite..... asp	Pyrite..... py
Lead..... Pb	Chalcopyrite..... cp	Pyrrhotite..... pyr
Molybdenum..... Mo	Fluorite..... fl	Quartz..... qtz
Nickel..... Ni	Galenite..... gal	Sphalerite..... sp

INDEX TO MINING PROPERTIES AND PROSPECTS

1. Atlantic Nickel (Rodgers Farm): cp, po, pn.
2. Clark Farm (A'Zone): cp, po.
3. Hall Carroll: po, cp.
4. Denis Stream (N'Zone): cp, po.
5. Grant Farm: cp, po.
6. 'C' Zone: cp, po.
7. Union Bridge - St. Croix River: cp, po.
8. Moores Mills - qtz, asp, gn, py.
9. Rollingdam Molybdenite: Mo, fl.
10. Mt. Blair: cp, fl, mal, hem.
11. Oliver - Cameron: qtz, cp, Bi, Au, Ag, mal.
12. T. Dick: qtz, py, cp, po, gn.
13. Lettice: qtz, cp, gn.
14. Simpson: sp, gn, py.
15. Daniel Hatt Farm: qtz, cp, po, gn.

Field work by W.J. Wolfe, M. Mason, G.J. Maserolle, J.M. Robbins, R.G. Garnett and D.K.J. MacKinnon
Analyses by Bondar-Clegg & Company, Limited, Ottawa, Ontario



Published, 1967
Copies of this map may be obtained from the
Mines Division, Department of Natural Resources,
Fredericton, New Brunswick